

Castelnuovo-Mumford regularity bound for smooth threefolds in $\mathbb{P}^5$ and extremal examples

By *Si-Jong Kwak* at KIAS

ABSTRACT. Let X be a nondegenerate integral subscheme of dimension n and degree d in $\mathbb{P}^N$ defined over the complex number field $\mathbb{C}$. X is said to be k -regular if $H^i(\mathbb{P}^N, \mathcal{I}_X(k-i)) = 0$ for all $i \geq 1$, where $\mathcal{I}_X$ is the sheaf of ideals of $\mathcal{O}_{\mathbb{P}^N}$ and Castelnuovo-Mumford regularity $\text{reg}(X)$ of X is defined as the least such k . There is a well-known conjecture concerning k -regularity: $\text{reg}(X) \leq \deg(X) - \text{codim}(X) + 1$. This regularity conjecture including the classification of borderline examples was verified for integral curves (Castelnuovo, Gruson, Lazarsfeld and Peskine), and an optimal bound was also obtained for smooth surfaces (Pinkham, Lazarsfeld). It will be shown here that $\text{reg}(X) \leq \deg(X) - 1$ for smooth threefolds X in $\mathbb{P}^5$ and that the only extremal cases are the Segre threefold and the complete intersection of two quadrics. Furthermore, every smooth threefold X in $\mathbb{P}^5$ is k -normal for all $k \geq \deg(X) - 4$, which is the optimal bound as the Palatini 3-fold of degree 7 shows. The same bound also holds for smooth regular surfaces in $\mathbb{P}^4$ other than for the Veronese surface.

0. Introduction.

Let X be a projective variety of dimension n and degree d in $\mathbb{P}^N$ defined over the complex number field $\mathbb{C}$. X is said to be k -normal if the homomorphism

$$H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}}(k)) \rightarrow H^0(X, \mathcal{O}_X(k))$$

is surjective, i.e., hypersurfaces of degree k cut out a complete linear system on X . We know that X is k -normal for all $k \gg 0$. It would be very interesting to find an *explicit* bound k_0 in terms of its invariants such that X is k -normal for all $k \geq k_0$ for all nondegenerate projective varieties.

On the other hand, as D.Mumford defined, we say that X is m -regular if $H^i(\mathbb{P}^N, \mathcal{I}_X(m-i)) = 0$ for all $i \geq 1$, where $\mathcal{I}_X$ is the sheaf of ideals of $\mathcal{O}_{\mathbb{P}^N}$. It is easy to check that X is $(m+1)$ -regular if and only if X is m -normal and $H^i(X, \mathcal{O}_X(m-i)) = 0$ for all $i > 0$. Let's define $\text{reg}(X)$ to be

This work is partially supported by KIAS and GARC(Global Analysis Research Center), Seoul, Korea

$\min\{m \in \mathbb{Z}: X \text{ is } m\text{-regular}\}$. More generally, for a coherent sheaf $\mathcal{F}$ on $\mathbb{P}^N$, $\mathcal{F}$ is m -regular if $H^i(\mathbb{P}^N, \mathcal{F}(m-i)) = 0$ for all $i \geq 1$ and $\text{reg}(\mathcal{F})$ is defined by $\inf\{m \in \mathbb{Z}: \mathcal{F} \text{ is } m\text{-regular}\}$.

We know that if X is m -regular, then the degrees of the minimal generators of the saturated ideal I_X defining X are bounded by m and hence there is no $(m+1)$ -secant line of X . (More generally, the degrees of generators of the i -th module of syzygies are bounded by $m+i$ [EG].)

Furthermore, the Hilbert polynomial and the Hilbert function of X have the same values for all $k \geq m-1$.

Regularity conjecture [EG], [GLP].

- (1) X is m -normal for all $m \geq d - \text{codim}(X)$.
- (2) X is m -regular for all $m \geq d - \text{codim}(X) + 1$, i.e.,
 $\text{reg}(X) \leq d - (N - n) + 1$.

In addition, the problem classifying all extremal examples which make the bound best possible has been one of interesting themes along these lines.

Historically, the curve case in $\mathbb{P}^3$ was first proved by Castelnuovo in 1893 [C] and his result was completed by Gruson, Lazarsfeld and Peskine in 1983 [GLP], where they proved, by using vector bundle techniques, that every integral curve X of degree d in $\mathbb{P}^N$ is $(d - \text{codim}(X) + 1)$ -regular. They also made the list of all extremal curves which fail to be $(d - \text{codim}(X))$ -regular

For smooth surfaces, bounds are obtained in [P] and [L], where R. Lazarsfeld got the optimal bound $\text{reg}(X) \leq (d - \text{codim}(X) + 1)$. For smooth threefolds and smooth fourfolds, bounds are obtained in [K], where the author got the following results: $\text{reg}(X) \leq (d - \text{codim}(X) + 2)$ for smooth threefolds and $\text{reg}(X) \leq (d - \text{codim}(X) + 5)$ for smooth fourfolds. Note that these bounds are off by 1 (resp., 4) from the conjectured bound. For arbitrary dimensional smooth projective varieties, $\text{reg}(X) \leq \min\{e, n+1\}(d-1) - n+1$, $e = \text{codim}(X)$ by Bertram, Ein and Lazarsfeld [BEL]. In the case of arbitrary codimension two smooth projective variety X of degree d in $\mathbb{P}^N$, Alzati [Al] got a better bound, i.e., $\text{reg}(X) \leq (d+2) + \frac{1}{2}N(N-1) - 2N$ (especially, $\text{reg}(X) \leq d + \frac{1}{2}N(N-1) - (N-1)[\frac{(N+4)}{2}]$, $N \geq 6$) by using various facts on higher order normality and vanishing theorems for an ideal sheaf due to Alzati and Ottaviani and recently, Peeva and Sturmfels showed in [PSt] that $\text{reg}(X) \leq \deg(X)$ for toric varieties of codimension two. However, unfortunately, for smooth projective varieties of dimension $n \geq 2$, extremal examples with geometric interpretations has not been classified yet.

The purpose of this paper is to give new bounds for regularity of smooth threefolds in $\mathbb{P}^5$ and to classify extremal examples by using local cohomology.

Theorem 1. *If a locally Cohen-Macaulay projective variety X of dimension n*

is contained in a hyperquadric Q in $\mathbb{P}^{n+2}$ then X is either a complete intersection or projectively Cohen-Macaulay and linked to a linear space $L \simeq \mathbb{P}^n$ via a complete intersection of Q and another hypersurface.

Theorem 2. *Let X be a smooth threefold of degree d in $\mathbb{P}^5$. Then $\text{reg}(X) \leq d - 1$. Furthermore, the only extremal examples are the Segre threefold and the complete intersection of two quadrics.*

Corollary 3. *Any smooth threefold in $\mathbb{P}^5$ is k -normal for all $k \geq d - 4$. The Palatini scroll of degree 7 which is not 2-normal (see example 2.0.) shows that this is the optimal bound. In addition, the same is true for regular surfaces in $\mathbb{P}^4$ except for the Veronese surface.*

Acknowledgements. I would like to express my gratitude to Professor Henry Pinkham for his encouragement and stimulating discussion with geometric insight throughout my graduate studies at Columbia University. In addition, I also thank my colleagues, especially Bo Ilic, for valuable discussions in the seminar group during my stay there.

1. Basic background. In this section, we recall the definitions and well-known results which are used in the following sections.

Lemma 1.1. *Let $\mathcal{F}$ be a p -regular vector bundle on $\mathbb{P}^n$ and $\mathcal{G}$ be a q -regular vector bundle on $\mathbb{P}^n$ defined over the complex numbers $\mathbb{C}$. Then $\mathcal{F} \otimes \mathcal{G}$ is $(p+q)$ -regular and $S^k(\mathcal{F})$ and $\bigwedge^k(\mathcal{F})$ are (kp) -regular.*

Proof. See [L], p.428.

Let X be a locally Noetherian scheme, Y a closed subscheme of X , and let $\mathcal{F}$ be a coherent sheaf on X . We define

$$\text{depth}_Y \mathcal{F} = \inf_{x \in Y} (\text{depth } \mathcal{F}_x),$$

where x is not necessarily a closed point.

Consider the following left exact functor $\underline{\Gamma}_Y(X, \cdot)$ from the category of sheaves of abelian groups to the category of sheaves of abelian groups, defined by

$$\underline{\Gamma}_Y(X, \mathcal{F})(U) = \Gamma_{Y \cap U}(U, \mathcal{F}|_U)$$

which is a subgroup of sections of $\mathcal{F}|_U$ with support in $Y \cap U$. Then the p -th right derived functors of $\underline{\Gamma}_Y(X, \cdot)$ are denoted by $\underline{H}_Y^p(X, \cdot)$, $p = 0, 1, \dots$. Similarly, consider the left exact functor $\Gamma_Y(X, \cdot)$ from the category of sheaves of abelian groups to the category of abelian groups, defined by $\mathcal{F} \mapsto \Gamma_Y(X, \mathcal{F})$ which is a subgroup of global sections of $\mathcal{F}$ with support in Y . We denote the p -th right derived functors of $\Gamma_Y(X, \cdot)$ by $H_Y^p(X, \cdot)$.

Note that $\underline{H}_Y^p(X, \cdot)$ is the sheaf associated to the presheaf

$$U \longmapsto H_{Y \cap U}^p(U, \mathcal{F}|_U)$$

and there is a spectral sequence $H_Y^n(X, \mathcal{F}) \Leftarrow E_2^{p,q} = H^p(X, \underline{H}_Y^q(\mathcal{F}))$.

Theorem 1.2. *Let X be a locally Noetherian scheme, Y a closed subscheme of X , and let $\mathcal{F}$ be a coherent sheaf on X . Then the following conditions are equivalent:*

- (1) $\underline{H}_Y^i(X, \mathcal{F}) = 0$ for all $i < n$,
- (2) $\text{depth}_Y \mathcal{F} \geq n$.

Proof. See [Gro] p.44.

We assume that X is a locally Cohen-Macaulay subscheme in $\mathbb{P}^N$ over the field $\mathbb{C}$ and $(M^i)(X) = H_*^i(\mathcal{I}_X) = \bigoplus_{m \in \mathbb{Z}} H^i(\mathbb{P}^N, \mathcal{I}_X(m))$ for $1 \leq i \leq \dim(X)$ and $[(M^i)(X)]^{\mathbb{C}} = \text{Hom}_{\mathbb{C}}((M^i)(X), \mathbb{C})$.

Theorem 1.3. *Let $X_1, X_2 \subset \mathbb{P}^N$ have dimension n . Assume that X_1 is linked to X_2 via a complete intersection $X = Z(F_1, F_2, \dots, F_{N-n})$ with $\deg(F_i) = d_i$. Let $d = \sum d_i$. Then*

- (1) $(M^{n-i+1})(X_2) \simeq [(M^i)(X_1)]^{\mathbb{C}}(N+1-d)$
- (2) *We have an exact sequence*

$$0 \rightarrow \mathcal{I}_X \rightarrow \mathcal{I}_{X_1} \rightarrow \omega_{X_2}(N+1-d) \rightarrow 0$$

where ω_{X_2} is the dualizing sheaf of X_2 .

Proof. See [Mi], Ch IV.

Theorem 1.4 (The (dimension+2)-secant Lemma). *Let $X \subset \mathbb{P}^N$ be a smooth n -dimensional subvariety and let Y be an irreducible variety parametrizing a family $\{L_y\}$ of lines in $\mathbb{P}^N$. Assume that, for a general L_y , the length of the scheme-theoretic intersection $L_y \cap X$ is at least $(n+2)$. Then we have*

$$\dim(\cup_{y \in Y} L_y) \leq n+1$$

Proof. See [Ran2].

2. Castelnuovo-Mumford Regularity for Smooth Threefolds in $\mathbb{P}^5$ and Extremal Examples.

Throughout this section, we work over the complex number field $\mathbb{C}$. A conjecture of R. Hartshorne (which is still open) states that any smooth n -manifold X in $\mathbb{P}^{n+2}$, ($n \geq 4$) is a complete intersection, equivalently it is projectively

normal. However, we have many examples of smooth non-projectively normal surfaces in $\mathbb{P}^4$ and 3-folds in $\mathbb{P}^5$, some of which are defined scheme-theoretically as the dependency loci of a morphism $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ between vector bundles $\mathcal{F}$ and $\mathcal{G}$ of rank m and $m + 1$ on $\mathbb{P}^N$, respectively. In other words, let $\mathcal{H}om(\mathcal{F}, \mathcal{G})$ be globally generated and φ be a generic morphism. Then the morphism φ induces

$$(\bigwedge^m(\mathcal{G}))^\vee \otimes (\det \mathcal{G}) \simeq \mathcal{G} \xrightarrow{(\bigwedge^m \varphi)^*} (\bigwedge^m(\mathcal{F}))^\vee \otimes (\det \mathcal{G}) \simeq \mathcal{O}_{\mathbb{P}^N}(c_1(\mathcal{G}) - c_1(\mathcal{F}))$$

which fits into an exact sequence

$$0 \rightarrow \mathcal{F} \xrightarrow{\varphi} \mathcal{G} \rightarrow \mathcal{I}_X(c_1(\mathcal{G}) - c_1(\mathcal{F})) \rightarrow 0,$$

where X is the dependency locus of $\varphi : \mathcal{F} \rightarrow \mathcal{G}$. By Kleiman's Bertini-type theorem [Klm], X is a codimension two subvariety in $\mathbb{P}^n$ and nonsingular away from a subset of codimension ≤ 4 in X .

Examples 2.0. From the exact sequence

$$0 \rightarrow \Omega_{\mathbb{P}^5}^2(2+t) \rightarrow \bigwedge^2(V) \otimes \mathcal{O}_{\mathbb{P}^5}(t) \rightarrow \Omega_{\mathbb{P}^5}(2+t) \rightarrow 0,$$

we know that $\Omega_{\mathbb{P}^5}(2+t)$ is globally generated for all $t \geq 0$. Then, four generic sections s_1, s_2, s_3, s_4 of $H^0(\mathbb{P}^5, \Omega_{\mathbb{P}^5}(2+t))$ induce an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^5}^{\oplus 4} \xrightarrow{\varphi=(s_1, s_2, s_3, s_4)} \Omega_{\mathbb{P}^5}^1(2+t) \xrightarrow{(\bigwedge^4 \varphi)^*} \mathcal{I}_{X_t}(c_1(\Omega_{\mathbb{P}^5}(2+t))) \rightarrow 0$$

where X_t is the dependency locus of φ . X_t is smooth by Kleiman's Bertini-type theorem [Klm]. By the formula [OSS] p.16,

$$c_k(E \otimes \mathcal{L}) = \sum_{i=0}^k \binom{r-i}{k-i} c_i(E) \cdot c_1(\mathcal{L})^{k-i}, \quad r = \text{rank}(E), \quad \mathcal{L} \text{ a line bundle},$$

we get $c_1(\Omega_{\mathbb{P}^5}^1(2+t)) = 4 + 5t$, $c_2(\Omega_{\mathbb{P}^5}^1(2+t)) = 10t^2 + 16t + 7 = \deg(X_t)$, and we can compute

$$h^1(\mathbb{P}^5, \mathcal{I}_{X_t}(k)) = \begin{cases} 1, & \text{when } k = 4t + 2 \\ 0, & \text{when } k \neq 4t + 2 \end{cases}$$

via the vanishing of $h^q(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^p(k))$ [OSS]. So, X_t is k -normal for all $k \neq 4t + 2$. Note that X_0 is called the Palatini scroll. It has degree 7 and sectional genus 4. According to a conjecture of Peskine and Van de Ven, it is the unique, non 2-normal, smooth threefold in $\mathbb{P}^5$.

Theorem 2.1. *If a locally Cohen-Macaulay, nondegenerate subvariety $X^n \subset \mathbb{P}^{n+2}$ ($n \geq 2$) is contained in a hyperquadric Q , then X is either a complete intersection or is projectively Cohen-Macaulay and linked to a linear space $L \simeq \mathbb{P}^n$ via a complete intersection of Q and another hypersurface.*

Proof. By a suitable change of coordinates, we may assume that Q is defined by a homogeneous polynomial $x_0^2 + x_1^2 + \cdots + x_k^2$ of rank $k + 1$. Let's consider all cases according to $\text{rank}(Q)$.

- Case I: $\text{rank}(Q) = k + 1 \geq 5$. In this case, X is a complete intersection. See [H2], exercise II.6.5(d), Klein's theorem.
- Case II: $\text{rank}(Q) = 4$. Then Q is the cone over a smooth quadric surface Q_1 in $\mathbb{P}^3$ with vertex $\Lambda \cong \mathbb{P}^{n-2}$. Let $\mathcal{U} = Q - \Lambda$. Then $X - \Lambda$ is a Cartier divisor of $\mathcal{U}$, so $\mathcal{I}_{X/Q}|_{\mathcal{U}}$ is a line bundle on $\mathcal{U}$. We have an isomorphism $\pi_{\Lambda}^* : Cl(Q_1) \xrightarrow{\sim} Cl(\mathcal{U}) = Cl(Q)$ where the projection $\pi_{\Lambda} : \mathcal{U} \rightarrow Q_1 \subset \mathbb{P}^3$ is an affine bundle with fibre $\mathbb{A}^{n-1}$ and $\text{Pic}(Q_1) = \text{Pic}(\mathcal{U}) \simeq \mathbb{Z} \oplus \mathbb{Z}$. We'll show that $\mathcal{I}_{X/Q}|_{\mathcal{U}}$ is of type (a,b), $|a - b| \leq 1$, where $\mathcal{I}_{X/Q}|_{\mathcal{U}} = \pi_{\Lambda}^*(\mathcal{O}_{Q_1}(a, b))$ for some $a, b \in \mathbb{Z}$. The proof of case II will be deferred until after some auxiliary propositions and lemmas.

Lemma 2.2. *Let $X^n \subset \mathbb{P}^{n+2}$, ($n \geq 2$) be a locally Cohen-Macaulay subvariety contained in a hyperquadric Q of rank 4. Then $\text{depth}_{\Lambda}(\mathcal{I}_{X/Q}) \geq 3$.*

Proof. Since $\Lambda^{n-2} \subset Q \subset \mathbb{P}^{n+2}$ and Q is locally Cohen-Macaulay,

$$\text{depth}_{\Lambda}(\mathcal{O}_Q) = \inf_{x \in \Lambda} \text{depth}(\mathcal{O}_{Q,x}) = \text{depth}(\mathcal{O}_{Q,\eta}) = \text{height } \eta = 3,$$

where η is the generic point of Λ^{n-2} . Since $n - 4 \leq \dim Z \leq n - 2$ for any irreducible component Z of $X \cap \Lambda$, we get $2 \leq \text{depth}(\mathcal{O}_{X,\wp}) = \dim(\mathcal{O}_{X,\wp}) \leq 4$ for any minimal prime ideal $\wp$ containing I_X and I_{Λ} . Therefore, $\text{depth}_{\Lambda}(\mathcal{O}_X) \geq 2$ whenever $X \cap \Lambda \neq \emptyset$. From a long exact sequence of local cohomology groups

$$\begin{aligned} (2.2.1) \quad \underline{H}_{\Lambda}^0(Q, \mathcal{I}_{X/Q}) &\rightarrow \underline{H}_{\Lambda}^0(Q, \mathcal{O}_Q) \rightarrow \underline{H}_{\Lambda}^0(Q, \mathcal{O}_X) \\ &\xrightarrow{\delta} \underline{H}_{\Lambda}^1(Q, \mathcal{I}_{X/Q}) \rightarrow \underline{H}_{\Lambda}^1(Q, \mathcal{O}_Q) \rightarrow \underline{H}_{\Lambda}^1(Q, \mathcal{O}_X) \\ &\rightarrow \underline{H}_{\Lambda}^2(Q, \mathcal{I}_{X/Q}) \rightarrow \underline{H}_{\Lambda}^2(Q, \mathcal{O}_Q) \rightarrow \underline{H}_{\Lambda}^2(Q, \mathcal{O}_X) \rightarrow \end{aligned}$$

Since $\text{depth}_{\Lambda}(\mathcal{O}_Q) = 3$ and $\text{depth}_{\Lambda}(\mathcal{O}_X) \geq 2$, by using Theorem 1.2.,

$$\text{depth}_{\Lambda}(\mathcal{I}_{X/Q}) \geq \min(\text{depth}_{\Lambda}(\mathcal{O}_Q), \text{depth}_{\Lambda}(\mathcal{O}_X) + 1) = 3.$$

□

Corollary 2.3. *We have isomorphisms $H^i(Q, \mathcal{I}_{X/Q}(m)) \simeq H^i(\mathcal{U}, \mathcal{I}_{X/Q}(m)|_{\mathcal{U}})$, for all $m \in \mathbb{Z}$ and $i = 0, 1$.*

Proof. For a coherent sheaf $\mathcal{I}_{X/Q}(m)$ on Q , we have a long exact sequence of cohomology groups

$$(2.3.1) \quad \begin{aligned} 0 \rightarrow H_{\Lambda}^0(Q, \mathcal{I}_{X/Q}(m)) &\rightarrow H^0(Q, \mathcal{I}_{X/Q}(m)) \rightarrow H^0(\mathcal{U}, \mathcal{I}_{X/Q}(m)|_{\mathcal{U}}) \\ &\rightarrow H_{\Lambda}^1(Q, \mathcal{I}_{X/Q}(m)) \rightarrow H^1(Q, \mathcal{I}_{X/Q}(m)) \rightarrow H^1(\mathcal{U}, \mathcal{I}_{X/Q}(m)|_{\mathcal{U}}) \\ &\rightarrow H_{\Lambda}^2(Q, \mathcal{I}_{X/Q}(m)) \rightarrow H^2(Q, \mathcal{I}_{X/Q}(m)) \rightarrow H^2(\mathcal{U}, \mathcal{I}_{X/Q}(m)|_{\mathcal{U}}) \end{aligned}$$

By Theorem 1.2 and Lemma 2.2, $H_{\Lambda}^i(Q, \mathcal{I}_{X/Q}(m)) = 0$ for $i = 0, 1, 2$. Therefore, we have isomorphisms $H^i(Q, \mathcal{I}_{X/Q}(m)) \simeq H^i(\mathcal{U}, \mathcal{I}_{X/Q}(m)|_{\mathcal{U}})$, $i = 0, 1$. $\square$

Lemma 2.4. *Let $Q = \text{Proj } \mathbb{C}[x_0, x_1, \dots, x_{n+2}]/(x_0x_3 - x_1x_2)$ be a hyperquadric of rank 4 in $\mathbb{P}^{n+2}$ with a vertex $\Lambda = Z(x_0, x_1, x_2, x_3)$ and $\Lambda = \mathbb{P}(V)$, $V = \mathbb{C}x_4 \oplus \mathbb{C}x_5 \oplus \dots \oplus \mathbb{C}x_{n+2}$. Let $\pi_{\Lambda} : \mathcal{U} = Q - \Lambda \rightarrow Q_1$ be a projection from the center Λ . Then $\pi_{\Lambda*} \mathcal{O}_{\mathcal{U}} = \bigoplus_{m=0}^{\infty} S^m V \otimes \mathcal{O}_{Q_1}(-m)$.*

Proof. Let $\{D^+(x_i) | i = 0, 1, 2, 3\}$ be the standard covering of Q_1 and $A_i = \mathbb{C}[\frac{x_0}{x_i}, \dots, \frac{x_3}{x_i}]/(\frac{x_0}{x_i} \frac{x_3}{x_i} - \frac{x_1}{x_i} \frac{x_2}{x_i})$ be the affine coordinate ring of $Q_1 \cap D^+(x_i)$ for $i = 0, 1, 2, 3$. Then

$$\pi_{\Lambda}|_{D^+(x_i)} : \text{Spec } A_i[\frac{x_4}{x_i}, \frac{x_5}{x_i}, \dots, \frac{x_{n+2}}{x_i}] \rightarrow \text{Spec } A_i$$

is a restriction of π_{Λ} . On $D^+(x_i) \cap D^+(x_j)$, $(\frac{x_k}{x_i})^m = (\frac{x_j}{x_i})^m \cdot (\frac{x_k}{x_j})^m$, $0 \leq i, j \leq 3$, and $4 \leq k \leq n+2$. Furthermore, $(\frac{x_j}{x_i})^m$ is the transition function of $\mathcal{O}_{Q_1}(-m)$. Therefore,

$$\pi_{\Lambda*} \mathcal{O}_{\mathcal{U}} = \bigoplus_{m=0}^{\infty} S^m(V) \otimes \mathcal{O}_{Q_1}(-m),$$

where $\Lambda = \mathbb{P}(V)$ and $S^m(V)$ is the m -th symmetric power of $V = \mathbb{C}x_4 \oplus \dots \oplus \mathbb{C}x_{n+2}$. This completes the proof of Lemma 2.4. $\square$

Proposition 2.5. *Under the same situation as Lemma 2.2,*

- (1) $|a - b| \leq 1$, where $\mathcal{I}_{X/Q}|_{\mathcal{U}} \simeq \pi_{\Lambda}^*(\mathcal{O}_{Q_1}(a, b))$
- (2) $H^1(Q, \mathcal{I}_{X/Q}(k)) = 0$ for all $k \in \mathbb{Z}$

Proof. Under the isomorphisms $\pi_{\Lambda}^* : Cl(Q_1) \xrightarrow{\sim} Cl(\mathcal{U}) = Cl(Q)$ and $Cl(Q_1) \simeq Cl(\mathbb{P}^1 \times \mathbb{P})$, we have $\mathcal{I}_{X/Q}|_{\mathcal{U}} = \pi_{\Lambda}^*(\mathcal{O}_{Q_1}(a, b))$ for some $a, b \in \mathbb{Z}$ and $\mathcal{I}_{X/Q}(k)|_{\mathcal{U}} = \pi_{\Lambda}^*(\mathcal{O}_{Q_1}(a+k, b+k))$. Since π_{Λ} is an affine morphism, so $R^i \pi_{\Lambda*}(\mathcal{O}_{\mathcal{U}}) = 0$, $i > 0$ and $H^i(\mathcal{U}, \mathcal{I}_{X/Q}(k)|_{\mathcal{U}}) = H^i(\mathcal{U}, \pi_{\Lambda}^* \mathcal{O}_{Q_1}(a+k, b+k)) = H^i(Q_1, \mathcal{O}_{Q_1}(a+k, b+k))$

$k) \otimes \pi_{\Lambda*} \mathcal{O}_{\mathcal{U}})$ for all $i \geq 0$. Therefore, for all $i = 0, 1$,

$$\begin{aligned} H^i(Q, \mathcal{I}_{X/Q}(k)) &\cong H^i(\mathcal{U}, \mathcal{I}_{X/Q}(k)|_{\mathcal{U}}) \\ &\cong H^i(Q_1, \mathcal{O}_{Q_1}(a+k, b+k) \otimes \pi_{\Lambda*} \mathcal{O}_{\mathcal{U}}) \\ &\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^i(Q_1, \mathcal{O}_{Q_1}(a+k-m, b+k-m)). \end{aligned}$$

Since $Q_1 \simeq \mathbb{P}^1 \times \mathbb{P}^1$, it follows immediately from the Künneth formula,

$$\begin{aligned} H^1(Q, \mathcal{I}_{X/Q}(k)) &\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^1(Q_1, \mathcal{O}_{Q_1}(a+k-m, b+k-m)) \\ &\cong \bigoplus_{m=0}^{\infty} \{S^m(V) \otimes H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(a+k-m) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(b+k-m))\} \oplus \\ &\quad \{S^m(V) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(a+k-m) \otimes H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(b+k-m))\} \end{aligned}$$

By Serre's vanishing Theorem, $H^1(Q, \mathcal{I}_{X/Q}(k)) = 0$ for $k \gg 0$. Therefore,

$$H^1(Q_1, \mathcal{O}_{Q_1}(a+k-m, b+k-m)) = 0, \text{ for all } m \in \mathbb{Z}^+ \text{ and for all } k \gg 0.$$

Let's fix an integer k_0 such that $H^1(Q, \mathcal{I}_{X/Q}(k_0)) = 0$ and assume $b \geq a$. We know that in general,

$$\begin{aligned} h^0(\mathcal{O}_{\mathbb{P}^1}(b+k_0-m)) &= h^0(\mathcal{O}_{\mathbb{P}^1}(a+k_0-m)) = 0, \text{ for all } m \geq b+k_0+1 \\ h^1(\mathcal{O}_{\mathbb{P}^1}(b+k_0-m)) &= h^1(\mathcal{O}_{\mathbb{P}^1}(a+k_0-m)) = 0, \text{ for all } m \leq a+k_0+1 \end{aligned}$$

If $a+2 \leq b$, then $\{m | a+k_0+2 \leq m \leq b+k_0\}$ is not empty. This implies that

$$H^1(Q_1, \mathcal{O}_{Q_1}(a+k_0-m, b+k_0-m)) \neq 0 \text{ for } m = a+k_0+2,$$

which contradicts the fact that $H^1(Q, \mathcal{I}_{X/Q}(k_0)) = 0$. So $|a-b| \leq 1$. In particular, by repeating this argument, $H^1(Q, \mathcal{I}_{X/Q}(k)) = 0$ for all $k \in \mathbb{Z}$. $\square$

Now let's go back to the proof of Case II in Theorem 2.1. Consider the exact sequence

$$0 \rightarrow \mathcal{I}_{Q/\mathbb{P}^{n+2}}(k) \rightarrow \mathcal{I}_{X/\mathbb{P}^{n+2}}(k) \rightarrow \mathcal{I}_{X/Q}(k) \rightarrow 0$$

Since $h^1(\mathbb{P}_{\mathbb{C}}^{n+2}, \mathcal{I}_{Q/\mathbb{P}^{n+2}}(k)) = h^1(Q, \mathcal{I}_{X/Q}(k)) = 0$ for all $k \in \mathbb{Z}$, we have $h^1(\mathcal{I}_{X/\mathbb{P}^{n+2}}(k)) = 0$ for all $k \in \mathbb{Z}$. Furthermore, if $\mathcal{I}_{X/Q}|_{\mathcal{U}} \cong \pi_{\Lambda}^* \mathcal{O}_{Q_1}(-a, -a)$ for

some $a > 0$, then $X - \Lambda$ is linearly equivalent to (a, a) -type in $Cl(\mathcal{U}) \cong \mathbb{Z} \oplus \mathbb{Z}$. So X is also linearly equivalent to a class of type (a, a) in $Cl(Q)$ and $\deg(X) = 2a$. On the other hand, we know that

$$\begin{aligned}
H^0(Q, \mathcal{I}_{X/Q}(a)) &\cong H^0(\mathcal{U}, \mathcal{I}_{X/Q}(a)|_{\mathcal{U}}) \\
&\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(Q_1, \mathcal{O}_{Q_1}(-m, -m)) \\
&\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-m)) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-m)) \\
&\cong \mathbb{C}
\end{aligned}$$

From the exact sequence

$$0 \rightarrow H^0(\mathbb{P}^{n+2}, \mathcal{O}_{\mathbb{P}^{n+2}}(a-2)) \xrightarrow{\times Q} H^0(\mathbb{P}^{n+2}, \mathcal{I}_{X/\mathbb{P}^{n+2}}(a)) \rightarrow H^0(Q, \mathcal{I}_{X/Q}(a)) \rightarrow 0$$

we get $h^0(\mathbb{P}^{n+2}, \mathcal{I}_{X/\mathbb{P}^{n+2}}(a)) = h^0(\mathbb{P}^{n+2}, \mathcal{O}_{\mathbb{P}^{n+2}}(a-2)) + 1$ and X is a complete intersection of Q and another hypersurface of degree a .

Similarly, if $\mathcal{I}_{X/Q}|_{\mathcal{U}} \cong \pi_{\Lambda}^* \mathcal{O}_{Q_1}(-a, -a+1)$ in $\text{Pic}(\mathcal{U})$, then X is linearly equivalent to $(a, a-1)$ -type in $Cl(Q)$ and $\deg(X) = 2a-1$. Note that

$$\begin{aligned}
H^0(Q, \mathcal{I}_{X/Q}(a)) &\cong H^0(\mathcal{U}, \mathcal{I}_{X/Q}(a)|_{\mathcal{U}}) \\
&\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(Q_1, \mathcal{O}_{Q_1}(-m, -m+1)) \\
(*) \quad &\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-m)) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-m+1)) \\
&\cong \mathbb{C} \oplus \mathbb{C}
\end{aligned}$$

and $h^0(\mathbb{P}^{n+2}, \mathcal{I}_{X/\mathbb{P}^{n+2}}(a)) = h^0(\mathbb{P}^{n+2}, \mathcal{O}_{\mathbb{P}^{n+2}}(a-2)) + 2$. Hence X is linked to $L \cong \mathbb{P}^n$ via a complete intersection of Q and another hypersurface of degree a . By the way, the property of being projectively Cohen-Macaulay is invariant in a liaison class by Theorem 1.3,(1). Thus X is projectively Cohen-Macaulay because L^n is projectively Cohen-Macaulay.

Case III: $\text{rank}(Q) = 3$. We may assume that $V = \mathbb{C}x_3 \oplus \mathbb{C}x_4 \oplus \cdots \oplus \mathbb{C}x_{n+2}$, $\text{Sing}(Q) = \Lambda^{n-1} = \mathbb{P}(V)$, $Q = \langle C, \Lambda \rangle$, where C is a smooth conic in $\mathbb{P}^2$. Since X is locally Cohen-Macaulay, we can get $\text{depth}_{\Lambda}(\mathcal{O}_Q) = 2$ and $\text{depth}_{\Lambda} \mathcal{O}_X \geq 1$. Hence, by (2.2.1), $\underline{H}_{\Lambda}^i(Q, \mathcal{I}_{X/Q}) = 0$ for $i = 0, 1$ and $H^0(Q, \mathcal{I}_{X/Q}(m)) \cong H^0(\mathcal{U}, \mathcal{I}_{X/Q}(m)|_{\mathcal{U}})$. Note that $Cl(Q) \cong Cl(\mathcal{U}) \cong Cl(C) \cong Cl(\mathbb{P}^1) \cong \mathbb{Z}$. If

$\mathcal{I}_{X/Q}|_{\mathcal{U}} = \pi_{\Lambda}^*(\mathcal{O}_C(-a)) = \pi_{\Lambda}^*(i^*(\mathcal{O}_{\mathbb{P}^1}(-2a)))$ for some $a > 0$, where $i : C \rightarrow \mathbb{P}^1$ is an isomorphism, we have $\deg(X) = 2a$ and

$$H^0(Q, \mathcal{I}_{X/Q}(a)) \cong H^0(\mathcal{U}, \mathcal{I}_{X/Q}(a)|_{\mathcal{U}}) \cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(C, \mathcal{O}_C(-m)) \cong \mathbb{C},$$

and the exact sequence

$$0 \rightarrow H^0(\mathbb{P}^{n+2}, \mathcal{O}_{\mathbb{P}^{n+2}}(a-2)) \xrightarrow{\times Q} H^0(\mathbb{P}^{n+2}, \mathcal{I}_{X/\mathbb{P}^{n+2}}(a)) \rightarrow H^0(Q, \mathcal{I}_{X/Q}(a)) \rightarrow 0$$

implies that X is a complete intersection of Q and another hypersurface F of degree a .

Next, if $\mathcal{I}_{X/Q}|_{\mathcal{U}} \cong \pi_{\Lambda}^*(i^*\mathcal{O}_{\mathbb{P}^1}(-2a+1))$, then $\deg(X) = 2a-1$ and

$$\begin{aligned} H^0(Q, \mathcal{I}_{X/Q}(a)) &\cong H^0(\mathcal{U}, \mathcal{I}_{X/Q}(a)|_{\mathcal{U}}) \\ &\cong H^0(C, i^*\mathcal{O}_{\mathbb{P}^1}(-2a+1) \otimes \mathcal{O}_C(a) \otimes \pi_{\Lambda*}\mathcal{O}_{\mathcal{U}}) \\ (**) \quad &\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(C, i^*\mathcal{O}_{\mathbb{P}^1}(-2a+1) \otimes \mathcal{O}_C(a-m)) \\ &\cong \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-2m+1)) \cong \mathbb{C} \oplus \mathbb{C} \end{aligned}$$

Similarly, $X + L^n$ is linearly equivalent to $Q \cdot F$, where F is a hypersurface of degree a . As in the case II, X is projectively Cohen-Macaulay. This completes the proof of Theorem 2.1. $\square$

Remark 2.6. For smooth surfaces in $\mathbb{P}^4$, Theorem 2.1 was also proved by A.Aure and L.Roth, see [Au] [Ro].

Corollary 2.7. *Let $X^n \subset \mathbb{P}^{n+2}$ ($n \geq 2$) be a locally Cohen-Macaulay subvariety contained in a hyperquadric Q of rank 3 (resp., 4) with $\text{Sing}(Q) = \Lambda$. Then,*

- (1) $\text{depth}_{\Lambda}(\mathcal{I}_{X/Q}) = 2$ (resp., 3)
- (2) *If X is not a complete intersection, then $\text{Sing}(Q) = \Lambda \subset X$.*

Proof. For a proof of (1), we first assume that $\text{rank}(Q) = 3$. Since X is projectively Cohen-Macaulay, $H^i(\mathbb{P}^{n+2}, \mathcal{I}_X) = H^i(Q, \mathcal{I}_{X/Q}) = 0$ for $i = 1, 2$. Therefore, by (2.3.1), for $k = 0$ or $k = 1$,

$$H^1(\mathcal{U}, \mathcal{I}_{X/Q}|_{\mathcal{U}}) \simeq \bigoplus_{m=0}^{\infty} S^m(V) \otimes H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-2m-2a+k)) \simeq H_{\Lambda}^2(Q, \mathcal{I}_{X/Q}),$$

which is obvious nonzero and so by Theorem 1.2., $\text{depth}_\Lambda(\mathcal{I}_{X/Q}) = 2$. Similarly, in the case of $\text{rank}(Q) = 4$, we have an exact sequence

$$0 \rightarrow H^2(\mathcal{U}, \mathcal{I}_{X/Q}|_{\mathcal{U}}) \rightarrow H_\Lambda^3(Q, \mathcal{I}_{X/Q}) \rightarrow H^3(Q, \mathcal{I}_{X/Q}) \rightarrow 0$$

Note that for surface X , $h^3(Q, \mathcal{I}_{X/Q}) = h^3(\mathbb{P}^4, \mathcal{I}_{X/\mathbb{P}^4}) = h^2(X, \mathcal{O}_X)$ may not be zero. By the way, $\text{depth}_\Lambda(\mathcal{I}_{X/Q}) = 3$ because $H^2(\mathcal{U}, \mathcal{I}_{X/Q}|_{\mathcal{U}}) \simeq \bigoplus_{m=0}^\infty S^m(V) \otimes H^2(Q_1, \mathcal{O}_{Q_1}(-m-a, -m-b))$, $|a-b| \leq 1$ is nonzero.

For (2), by Theorem 2.1, we know that $X = Q \cap F_1 \cap F_2$, where F_i is a homogeneous polynomial of degree a , $\deg(X) = 2a - 1$ and $Q \cap F_i = X \cup L_i$, $L_i = \langle \Lambda, \ell_i \rangle$ where ℓ_i is a line in $Q_1 \subset \mathbb{P}^3$ (or ℓ_i is one point of a smooth conic C in $\mathbb{P}^2$ if $\text{rank}(Q) = 3$). Thus, $\Lambda \subset X = Q \cap F_1 \cap F_2$ $\square$

Remark 2.8. It is well-known that if $X^n \subset \mathbb{P}^{n+2}$, $n \geq 5$ and X a locally Cohen-Macaulay variety, then X is a complete intersection if and only if X is projectively Cohen-Macaulay, see [PSz].

Proposition 2.9. *If a locally Cohen-Macaulay variety $X^n \subset \mathbb{P}^{n+2}$, ($n \geq 2$) is contained in a hyperquadric Q and is not a complete intersection, then*

- (1) *The saturated ideal $I_X = \bigoplus_{m \in \mathbb{Z}} H^0(\mathbb{P}^{n+2}, \mathcal{I}_X(m))$ is generated by polynomials Q, F_1, F_2 , $\deg(F_i) = a$ and $\deg(X) = 2a - 1$.*
- (2) *$\text{reg}(X) = \frac{\deg(X)+1}{2} = a$.*

Proof. Since X is not a complete intersection, $\text{rank}(Q) = 3$ or 4 , as we have already noted. Assume that $\mathcal{I}_{X/Q}|_{\mathcal{U}} = \pi_\Lambda^*(\mathcal{O}_{Q_1}(-a, -a+1))$ or $\pi_\Lambda^*(i^*\mathcal{O}_{\mathbb{P}^1}(-2a+1))$ as in the proof of Theorem 2.1. By (*) and (**), we can show that for $a \geq 2$

$$h^0(Q, \mathcal{I}_{X/Q}(k)) = \begin{cases} 0 & \text{when } 2 \leq k < a, \\ 2 & \text{when } k = a. \end{cases}$$

Let $\alpha(I_X) = \min\{i \in \mathbb{Z} : h^0(\mathcal{I}_X(i)) \neq 0\} = 2$ and $\nu(I_X)$ = number of minimal generators of the saturated ideal I_X of X . Since X is projectively Cohen-Macaulay of codimension two, we have $\nu(I_X) = \alpha(I_X) + 1 = 3$ (see [Mi], Corollary 2.2.3.). Therefore, the saturated ideal $I_X = \bigoplus_{m \in \mathbb{Z}} H^0(\mathbb{P}^{n+2}, \mathcal{I}_X(m))$ is generated by polynomials Q, F_1, F_2 , $\deg(F_i) = a$. For a proof of (2), from (*) and (**), we have $h^0(Q, \mathcal{I}_{X/Q}(a+1)) = 2n + 4$. The vector space of all polynomials of the form $c_1 F_1 \cdot L_1 + c_2 F_2 \cdot L_2$, where L_1 and L_2 are arbitrary linear forms, has dimension $(2n + 6)$. This implies that there are two relations among minimal generators Q, F_1, F_2 in degree $(a + 1)$. Thus, the minimal free resolution for I_X is a short exact sequence as follows:

$$0 \rightarrow S(-a-1) \oplus S(-a-1) \rightarrow S(-2) \oplus S(-a) \oplus S(-a) \rightarrow I_X \rightarrow 0.$$

Consequently, $\text{reg}(X) = \frac{\deg(X)+1}{2} = a$. $\square$

Remark 2.10. If $X \subset \mathbb{P}^N$ is a complete intersection defined by polynomials $F_1, F_2, \dots, F_e$, $\deg(F_i) = d_i$, $e = \text{codim}(X, \mathbb{P}^N)$, then it is easily checked that $\text{reg}(X) = d_1 + d_2 + \dots + d_e - e + 1$.

Theorem 2.11. *Let X be a smooth threefold in $\mathbb{P}^5$ which is not contained in any hyperquadric Q . Then $\text{reg}(X) \leq \deg(X) - 3$. We also have $\text{reg}(S) \leq \deg(S) - 3$ for any regular surface S with $h^0(\mathcal{I}_{S/\mathbb{P}^4}(2)) = 0$ in $\mathbb{P}^4$ except for the Veronese surface.*

Proof. The following construction and technique are basically due to Lazarsfeld [L] and Alzati [Al]. The author wants to give more simplified and self-contained proof. Let $\mathbb{P}^5 = \text{Proj } \mathbb{C}[T_0, T_1, T_2, T_3, T_4, T_5]$. Consider the following diagram

$$\begin{array}{ccc} Bl_p(\mathbb{P}^5) = \mathbb{P}(\mathcal{O}_{\mathbb{P}^4}(1) \oplus \mathcal{O}_{\mathbb{P}^4}) & \xrightarrow{\pi_2} & \mathbb{P}^4 \\ \pi_1 \downarrow & & \\ X \subset \mathbb{P}^5 & & \end{array}$$

where $p = (0, 0, 0, 0, 0, 1) \notin X$ and $\pi_p : \mathbb{P}^5 \dashrightarrow \mathbb{P}^4$ is a generic projection, which commutes with the above diagram, i.e. $\pi_2 \circ \pi_1^{-1}|_X = \pi_p|_X$. Therefore, the natural morphism $\mathcal{O}_{\mathbb{P}^5}(k) \rightarrow \mathcal{O}_X(k)$ induces a morphism $w_k : \pi_{2*}(\pi_1^* \mathcal{O}_{\mathbb{P}^5}(k)) \cong \text{Sym}^k(\mathcal{O}_{\mathbb{P}^4}(1) \oplus \mathcal{O}_{\mathbb{P}^4}) \rightarrow \pi_{2*}(\pi_1^* \mathcal{O}_X(k)) \simeq \pi_{p*} \mathcal{O}_X(k)$.

By Theorem 1.4 (the (dimension+2)-secant Lemma), the length of each fibre of π_p is at most 4. Let L_y be the line through p, y . Then

$$w_3 \otimes \mathbb{C}(y) : H^0(L_y, \mathcal{O}_{L_y}(3)) \rightarrow H^0(L_y, \mathcal{O}_{\pi_p^{-1}(y)}(3))$$

is surjective, i.e., the scheme-theoretic fibre $\pi_p^{-1}(y)$ is 3-normal for all $y \in \pi_p(X)$ and by tensoring $\mathcal{O}_{\mathbb{P}^4}(-3)$, we have an exact sequence,

$$0 \rightarrow E \rightarrow \mathcal{O}_{\mathbb{P}^4}(-3) \oplus \mathcal{O}_{\mathbb{P}^4}(-2) \oplus \mathcal{O}_{\mathbb{P}^4}(-1) \oplus \mathcal{O}_{\mathbb{P}^4} \xrightarrow{w_3 \otimes \mathcal{O}_{\mathbb{P}^4}(-3)} \pi_{p*} \mathcal{O}_X \rightarrow 0$$

where $E = \text{Ker } w_3 \otimes \mathcal{O}_{\mathbb{P}^4}(-3)$.

Lemma 2.12. *The vector bundle $E = \text{Ker } w_3 \otimes \mathcal{O}_{\mathbb{P}^4}(-3)$ controls the regularity of X , i.e., $\text{reg}(X) \leq \text{reg}(E)$.*

Proof. According to Lazarsfeld [L], this is due to Gruson and Peskine. Suppose that E is $(m+1)$ -regular: in other words, $H^i(\mathbb{P}^4, E(m+1-i)) = 0$, $i > 0$. We'll show that X is $(m+1)$ -regular, equivalently, X is m -normal and $H^i(X, \mathcal{O}_X(m-i)) = 0$, $i \geq 1$. Consider the sequence

$$0 \rightarrow E \rightarrow \mathcal{O}_{\mathbb{P}^4}(-3) \oplus \mathcal{O}_{\mathbb{P}^4}(-2) \oplus \mathcal{O}_{\mathbb{P}^4}(-1) \oplus \mathcal{O}_{\mathbb{P}^4} \xrightarrow{w_3 \otimes \mathcal{O}_{\mathbb{P}^4}(-3)} \pi_{p*} \mathcal{O}_X \rightarrow 0$$

Since $H^1(\mathbb{P}^4, E(m)) = 0$, the following morphism

$$\begin{array}{c} H^0(\mathcal{O}_{\mathbb{P}^4}(m-3)) \oplus H^0(\mathcal{O}_{\mathbb{P}^4}(m-2)) \oplus H^0(\mathcal{O}_{\mathbb{P}^4}(m-1)) \oplus H^0(\mathcal{O}_{\mathbb{P}^4}(m)) \\ \downarrow H^0(w_3) \\ H^0(\pi_{p*}\mathcal{O}_X(m)) \end{array}$$

is surjective, where $H^0(\pi_{p*}\mathcal{O}_X(m)) \simeq H^0(X, \mathcal{O}_X(m))$ and $H^0(w_3)$ maps $F_i \in H^0(\mathcal{O}_{\mathbb{P}^4}(m-i))$ to $F_i \cdot T_5^i$ for $i = 0, 1, 2, 3$. Hence X is m -normal.

Secondly, $H^i(\mathbb{P}^4, E(m+1-i)) = 0$, $i \geq 2$ implies $H^i(X, \mathcal{O}_X(m-i)) = 0$ for $i \geq 1$. Indeed, from the exact sequence

$$0 \rightarrow E \rightarrow \mathcal{O}_{\mathbb{P}^4}(-3) \oplus \mathcal{O}_{\mathbb{P}^4}(-2) \oplus \mathcal{O}_{\mathbb{P}^4}(-1) \oplus \mathcal{O}_{\mathbb{P}^4} \xrightarrow{w_3} \pi_{p*}\mathcal{O}_X \rightarrow 0$$

we get, for $i \geq 1$,

$$H^i(X, \mathcal{O}_X(m-i)) = H^i(\mathbb{P}^4, \pi_{p*}\mathcal{O}_X(m-i)) = H^{i+1}(\mathbb{P}^4, E(m-i)) = 0$$

Therefore, X is $(m+1)$ -regular, and $\text{reg}(X) \leq \text{reg}(E)$. $\square$

Claim. E^* is (-3) -regular for a smooth 3-fold X in $\mathbb{P}^5$.

Proof of Claim. Note that E^* is (-3) -regular if and only if $H^i(\mathbb{P}^4, E^*(-3-i)) = 0$, $i = 1, 2, 3, 4$. For $i = 1$, $h^1(E^*(-4)) = h^3(E(-1)) = h^2(\pi_{p*}\mathcal{O}_X(-1)) = h^2(\mathcal{O}_X(-1)) = 0$ by Kodaira's vanishing theorem. When $i = 2$, $h^2(\mathbb{P}^4, E^*(-5)) = h^2(E) = h^1(\pi_{p*}\mathcal{O}_X) = h^1(\mathcal{O}_X) = 0$ by the Barth-Larsen theorem [B]. For $i = 3$, $h^3(\mathbb{P}^4, E^*(-6)) = h^1(E(1)) = 0$ because we have the following exact sequence

$$H^0(\mathcal{O}_{\mathbb{P}^4}(-2) \oplus \mathcal{O}_{\mathbb{P}^4}(-1) \oplus \mathcal{O}_{\mathbb{P}^4} \oplus \mathcal{O}_{\mathbb{P}^4}(1)) \xrightarrow{\alpha} H^0(\mathcal{O}_X(1)) \rightarrow H^1(E(1)) \rightarrow 0$$

and α is surjective by Zak's linear normality theorem [Z]. Finally, if $i = 4$, then $h^4(E^*(-7)) = h^0(E(2)) = 0$ because X is not contained in any hyperquadric and hence

$$H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(-1) \oplus \mathcal{O}_{\mathbb{P}^4} \oplus \mathcal{O}_{\mathbb{P}^4}(1) \oplus \mathcal{O}_{\mathbb{P}^4}(2)) \rightarrow H^0(\mathbb{P}^4, \pi_{p*}\mathcal{O}_X(2))$$

is injective. Therefore, E^* is (-3) -regular. This completes the proof of our claim. $\square$

Now, let's go back to the proof of Theorem 2.11. Note that $E \simeq \wedge^3 E^* \otimes \det E$, $\text{rank}(E) = 4$. So, by Lemma 1.1, $\text{reg}(E) \leq \text{reg}(\wedge^3(E^*)) + \text{reg}(\det(E)) \leq d-3$ where $\det(E) = \mathcal{O}_{\mathbb{P}^4}(-6-d)$ and $\text{reg}(\det(E)) = d+6$. Thus

$$\text{reg}(\mathcal{I}_X) = \text{reg}(X) \leq \text{reg}(E) \leq d-3$$

Furthermore, suppose that S is a smooth regular surface in $\mathbb{P}^4$ with degree d and $h^0(\mathcal{I}_S(2)) = 0$. By the same generic projection method, we have an exact sequence

$$0 \rightarrow E \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2) \oplus \mathcal{O}_{\mathbb{P}^3}(-1) \oplus \mathcal{O}_{\mathbb{P}^3} \xrightarrow{w_2 \otimes \mathcal{O}_{\mathbb{P}^3}(-2)} \pi_{p*} \mathcal{O}_S \rightarrow 0$$

Since S is linearly normal and $h^1(\mathcal{O}_S) = 0$, by a similar method, E^* is (-3) -regular and $\text{reg}(S) \leq \text{reg}(E) = \text{reg}(\bigwedge^2(E^*) \otimes \det(E)) \leq d - 3 \quad \square$

Remark 2.13.

- (a) Such a bound $(d - 3)$ in Theorem 2.11 is the best possible because, as shown in example 2.0., the dependency locus X_0 of four generic sections of $\Omega_{\mathbb{P}^5}(2)$ has degree 7 and $\text{reg}(X) = 4$. Such a threefold is called the Palatini scroll of degree 7 which is not 3-regular.
- (b) By Proposition 2.9 and Theorem 2.11, for a smooth 3-fold X in $\mathbb{P}^5$, $\text{reg}(X) \leq d - 1$ and $\text{reg}(X) = d - 1$ if and only if X has degree 3 or 4, i.e., it is either the Segre threefold or the complete intersection of two quadrics.
- (c) Theorem 2.11 is also true for smooth 3-folds in $\mathbb{P}^5$ defined over any algebraically closed field of characteristic zero because Kodaira's vanishing theorem can be extended to this case.

Corollary 2.14. *Let X be a smooth 3-fold in $\mathbb{P}^5$ of degree d . Then X is k -normal for $k \geq d - 4$. The Palatini scroll of degree 7 which, as we have already noted, is not 2-normal, shows that this is the optimal bound.*

Proof. By Theorem 2.1 and Theorem 2.11, this is clear. $\square$

REFERENCES

- [Al] A. Alzati, *A new Castelnuovo bound for two codimensional subvarieties of $\mathbb{P}^r$* , Proc. Amer. Math. Soc. **114** (1992), 607–611.
- [Au] A. Aure, *On surfaces in projective 4-space*, Thesis, Oslo (1987).
- [B] W. Barth, *Transplanting cohomology classes in complex projective space*, Amer. J. Math. **114** (1970), 951–967.
- [BEL] A. Bertram, L. Ein, and R. Lazarsfeld, *Vanishing theorems, a theorem of Severi, and the equations defining projective varieties*, J. of Amer. Math. Soc. **4** (1991), 587–602.
- [BM] D. Bayer and D. Mumford, *Proceedings of Computer Algebra as a Tool For Research in Mathematics and Physics*, Courant Institute, April, 1984.
- [BSS] M. Beltrametti, M. Schneider, A. Sommese, *Threefolds of degree 9 and 10 in $\mathbb{P}^5$* , Math. Ann. **288** (1990), 413–444.
- [C] G. Castelnuovo, *Sui multipli di una serie lineare di gruppi di punti appartenente ad una curva algebrica*, Rend. Circ. Mat. Palermo (2) **7** (1893), 89–110.
- [EG] D. Eisenbud and S. Goto, *Linear free resolutions and minimal multiplicity*, J. Alg. **88** (1984), 89–133.

- [G] V. A. Greenberg, *A Castelnuovo bound for projective varieties admitting a stable linear projection onto a hypersurface*, Ph.D. Thesis, Columbia University.
- [Gro] A. Grothendieck, *Local Cohomology (Notes by R. Hartshorne)*, Lecture Notes in Math. 41, Springer Verlag, Berlin-Heidelberg-New York, 1967.
- [GLP] L. Gruson, R. Lazarsfeld, and C. Peskine, *On a theorem of Castelnuovo and the equations defining projective varieties*, Inv. Math. **72** (1983), 491–506.
- [H1] R. Hartshorne, *Varieties of Small Codimension in Projective Space*, Bull. Amer. Math. Soc. **80** (1974), 1017–1032.
- [H2] ———, *Algebraic Geometry*, Springer Verlag, Heidelberg, 1977.
- [Ilic] B. Ilic, *Geometric properties of the double-point divisor*, Thesis (1995), Columbia University.
- [K] S. Kwak, *Castelnuovo Regularity of Smooth Projective Varieties of Dimension 3 and 4*, To appear in the J. of Algebraic Geometry.
- [Klm] S. Kleiman, *Geometry on grassmanians and applications to splitting bundles and smoothing cycles*, Publ. Math. I.E.H.S. **36** (1969), 281–297.
- [L] R. Lazarsfeld, *A Sharp Castelnuovo Bound for Smooth Surfaces*, Duke Math. J. **55** (1987), 423–438.
- [Mi] J. C. Migliore, *An Introduction to deficiency module and liaison theory for subschemes of projective space*, Lecture Note Series, No.24, Global Analysis Research Center, Research Institute of Mathematics, Seoul, Korea, 1994.
- [Mu] D. Mumford, *Lectures on curves on an algebraic surfaces*, Annals of Math. Studies, No.59, 1966.
- [OSS] Ch. Okonek, H. Spindler, and M. Schneider, *Vector bundles on complex projective spaces*, Birkhäuser, 1980.
- [P] H. Pinkham, *A Castelnuovo bound for smooth surfaces*, Inv. Math. **83** (1986), 321–332.
- [PSt] I. Peeva and B. Sturmfels, *Syzygies of codimension 2 lattice ideals*, preprint.
- [PSz] Ch. Peskine and L. Szpiro, *Liaison des varietes algebriques I*, Invent. Math. **26** (1974), 271–302.
- [Ran1] Z. Ran, *Local differential geometry and generic projections of threefolds*, J. Diff. Geo. **32** (1990), 131–137.
- [Ran2] ———, *The (dimension+2) secant lemma*, Inv. Math. **106** (1991), 65–71.
- [Rao] P. Rao, *Liaison Equivalence Classes*, Math. Ann. **258** (1981), 169–173.
- [Ro] L. Roth, *On the Projective Classification of Surfaces*, Proc. of London Math. Soc. **42** (1937), 142–170.
- [Se] F. Severi, *Intorno ai punti tripli apparenti.*, Rend. Circ. Mat. Palermo **15** (1901), 33–51.
- [Zak] F. Zak, *Tangents and secants of algebraic varieties*, Translation of Math. Monographs, vol.127, American Math. Society, 1993.